Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME 34

New names, and the final members of new combinations, are in **bold face type.**

Abies Douglasii, 591
Abrams, L. R. Studies on the flora of Southern California, 263
Acanthorhynchus, 313; **Vaccinii,** 314
Acanthospermum australe, 368
Acer, 284; dasycarpum, 369; platanoides, 234; rubrum, 367; saccharinum, 369
Acerates, 205
Acnida cannabina, 79; tamariscina, 79; tamariscina tuberculata, 78
Acrocryphaea Coffeae, 573
Acropterygium, 250, 251
Acrostichum furcatum, 243, 257
Adenostegia **ciliosa,** 35; ramosa, 35
Adenostoma fasciculatum, 264; fasciculatum densifolium, 263
Aecidium Amorphae, 590; **Boehmeriae,** 590; candidum, 64; **Cannonii,** 210; **fluxum,** 590; Ipomoeae, 69; Ipomoeae-panduranae, 68; **Petalostemonis,** 589; Sarcobati, 210; Urticae, 591
Aeschynomene virginica, 364
Aesculus glabra, 349; Pavia, 352
Agaricus amboinensis, 477; bulbosus, 348; campestris, 302; **Pattersonae,** 347
Ageratum conyzoides, 74
Agrostis, 585, 607; hyemalis, 584; perennans, 584; Thurberiana, 585
Albugo, 61, 63, 65, 73, 77, 82, 84, 387; Bliti, 62, 64, 76, 77, 79, 81, 82; brasiliensis, 74; candida, 61, 63, 64, 68, 69, 77, 81; Cruciferarum, 63; Ipomoeae-panduranae, 61, 64, 68, 81; Lepigoni, 61, 64, 71, 81; **occidentalis,** 62, 64, 80, 81; platensis, 62, 64, 77, 79, 81, 82; Portulacae, 62, 64, 77, 81, 83; quadrata, 64, 76; **sibirica,** 61, 63, 68; Solivae, 74; spinulosa, 73; **Swertiae,** 62, 64, 72, 81; **Tillaeae,** 64, 76; Tragopogonis, 62, 64, 70, 72, 74, 76, 81; **tropica,** 61–63, 68, 81, 82
Albugo, The genus, 61
Alnus rugosa, 367
Alopecurus, 398
Alyssum, 183; macrocarpum, 181
Amaranthus, 77, 82; Bigelovii, 78; blitoides, 78; Blitum, 77, 79; crispus, 79; emarginatus, 78; graecizans, 78; hybridus, 78, 81; hybridus paniculatus, 79; Palmeri, 78; retroflexus, 78; spinosus, 78; tristis, 79; viridis, 79
Amauroderma **Elmerianum,** 475
Ambrosia artemisiaefolia, 75, 364, 404; psilostachya, 75, 404; trifida, 75, 404
Amelanchier, 461; alnifolia, 462, 463; oreophila, 463
America, Mosses from tropical, 569
American botanical literature, Index to (1901–1905), 51; (1906), 105, 267, 575; (1907), 161, 213, 319, 379, 439, 483, 521, 609
American Code of Botanical Nomenclature, 167
American ferns, 243
American fossil mosses, 139
American species of Antrophyum, 445
Amorpha canescens, 590; herbacea, 365
Ampelopsis quinquefolia, 408; Ritchii, 409; Veitchii, 409
Anchistea virginica, 367
Andromeda floribunda, 352; **grandifolia,** 204, 206; latifolia, 204; Novae-Caesareae, 204; Parlatorei, 188, 203, 204, 206
Andropogon tener, 373
Androsace, 517–520; arizonica, 519; diffusa, 517–520; **glandulosa,** 518, 519; occidentalis, 517–520; pinetorum, 517–519; **platysepala,** 518, 519; puberulenta, 518, 519; septentrionalis, 517, 518; subulifera, 518
Androsace in New Mexico, 517
Antennaria plantaginifolia, 76
Anthostomella, 313; **destruens,** 313; Smilacis, 313
Antrophyopsis, 447
Antrophyum, 445–458; anetioides, 449, 455; Boryanum, 447; brasilianum, 448, 449, 452, 454–456; carnosum, 449; cayennense, 448, 449, 453; Desvauxii, 457; discoideum, 448, 449, 454, 455; **Dussianum,** 449, 453; ensiforme, 448–450; falcatum, 449; Feei, 451, 452; Galeottii, 449; giganteum, 457; immersum, 447; **Jenmani,** 449, 453, 454; lanceolatum, 445, 448, 449, 451, 452, 454; lineatum, 448–452; minimum, 457; reticulatum, 445; spathulatum, 452, 453; subsessile, 448, 452–455, 457; Werckleanum, 458

Antrophyum, The genus, 445
Apeibopsis discolor, 199
Arabis, 429; Drummondii alpina, 437; furcata, 65; glabra, 68; longirostris, 429; Lyallii, 437; lyrata, 65; **oreophila,** 437; rostrata, 429; virginica, 65
Aragallus, 423; **angustatus,** *422*; argophyllus, 423; **atropurpureus,** 424; Besseyi, 422–424; Blankinshipii, 422–424; collinus, 423, 424; Lagopus, 423; Lambertii, 421–424; monticolus, 424; multiceps, 423; nanus, 423; **patens,** 421; sericeus, 421, 424
Aralia, *201*; groenlandica, *201*; **Newberryi,** *201*, *206*; palmata, *201*; spinosa, 352, 366
Arenaria marina, *72*
Aristida stricta, 351, 365
Armillaria, 104; **ventricosa,** 104
Aronia arbutifolia, 365
Artemisia biennis, 75; canadensis, 76; ludoviciana, 405
ARTHUR, J. C. New species of Uredineae, 583
Asclepias, *221*; Cornuti, *239*; lanceolata, 364
Asimina triloba, 103
Aspergillus niger, 303
Aster Novae-Angliae, 394; oblongifolius, 394; sagittifolius, 394; squarrosus, 365
Astragalus, 49; atratus, 48; campestris, 49, 50; convallarius, 49; divergens, 417; Haydenianus, 420; multiflorus, 419; tegetarius, 421
Atlantic coastal plain, Mesozoic flora of the, 185
Atriplex, 82
Avena sativa, 286, 287
Avrainvillea, 491, 505–508, 512, 513; laetivirens, 512; levis, 508, 512; longicaulis, 504, 508–512; Mazei, 504, 509, 511, 512; nigricans, 504, 505, 508–512, 515; **Rawsoni,** 508, 510, 511, 516; sordida, 512
Avrainvillea, Further notes on Halimeda and, 491
Aytonia **Evansii,** 57–59; **jamaicensis,** 58, 60
Aytonia from Jamaica, Two new species of, 57
Azolla caroliniana, 364

Barbarea Barbarea, 65
BARNHART, J. H. A new Utricularia from Long Island, 579
Basidiophora, 388, 393; entospora, 393; **Kellermanii,** 393, 394, 405
Batodendron arboreum, 364
Bazzania, 561
BENEDICT, R. C. The genus Antrophyum, 445
BERRY, E. W. Contributions to the Mesozoic flora of the Atlantic coastal plain, 185
Beta, 82
Betula atavina, 193; nigra, 367
Bidens cernua, 404; comosa, 405; connata, 405; frondosa, 404; laevis, 404
Bignonia crucigera, 364
Blitum, 82; capitatum, 81, 82
Boehmeria cylindrica, 591
Boerhaavia anisophylla, 80; decumbens, 80, 81; diffusa, 80; erecta, 80; hirsuta, 80; paniculata, 80; Sonorae, 80; spicata, 80; viscosa, 80; Xanti, 80
Boletus, 86; pavonius, 468
Bonamia, 143, 146, 147; alternifolia, 146; aquatica, 149; grandiflora, 146; humistrata, 148; ovalifolia, 146; Pickeringii, 149
Botanical literature, American (1901–1905), 51; (1906), 105, 267, 575; (1907), 161, 213, 319, 379, 439, 483, 521, 609
Botanical Nomenclature, American Code of, 167
Bothrodiscus, 312; **pinicola,** 313
Botrytis, 498; cana, 408; intestans, 391, 408; macrospora, 401; nivea, 401; viticola, 407, 408; vitis-viticola, 407
Bouteloua oligostachya, *208*; prostrata, ***208***
Brachysteleum, 141
Bradburya virginiana, 364
Brassica, 66; arvensis, 66; campestris, 66; integrifolia, 66; Napus, 68; nigra, 65, 66; oleracea, 68
Breweria, 146, 147; angustifolia, 148; aquatica, 149; Choisyana, 148; grandiflora, 146; humistrata, 148; mexicana, 144; ovalifolia, 146; Pickeringii, 149; rotundifolia, 149; tenella, 148; trichosanthes, 148; villosa, 149
BRITTON, E. G. & HOLLICK, A. American fossil mosses, with description of a new species from Florissant, Colorado, 139
Bruguiera caryophylloides, *272*; eriopetala, *272*
Bryolejeunea, 558
Bryonia, *222*, 330, 339, 340; alba, *222*; dioica, *222*, 331
Bryonopsis lacunosa erythrocarpa, 414
Bryopteris, 4, 533, 557–559, 561, 564, 566; diffusa, 557–559, 561; filicina, 557–559, 563–566, 568; fruticulosa, 565; Gaudichaudii, 558; tenuicaulis, 559, 565, 566; trinitensis, 566; trinitensis intermedia, 566
Bryum, 571; argenteum, 570

BURLINGHAM, G. S. Some Lactarii from Windham County, Vermont, 85
Bursa, 65; Bursa-pastoris, 65, 66, 68

Caeoma Amaranthi, 77; convolvulatum, 68; **occidentale,** 591
Cakile edentula, 68; maritima, 66
California, Studies on the flora of Southern, 263
Calliospora **Petalostemonis,** 588
Callistolejeunea, 553
Calonyction aculeatum, 70
Calycobolus, 143; **amazonicus,** 145; emarginatus, 144; **ferrugineus,** 145; **glaber,** 145; **Pringlei,** 144, 145; pulchellus, 145; **sericeus,** 144, 145; **spectabilis,** 146; **umbellatus,** 146; **velutinus,** 144, 145
Camelina microcarpa, 65, 66; sativa, 66
Campulosus aromaticus, 363, 365
Campylopus concolor, 569; **Pittieri,** 569
Canadian Rocky Mountains, Two new willows from the, 157
Capnoides **brachycarpum,** 426; Cusickii, 426; **hastatum,** 426
Capsella Bursa-pastoris, 181, 182
Carallia integerrima, 272
Cardamine bulbosa, 66
Carduus, 73; arvensis, 75; lanceolatus, 75; muticus, 75; spinosissimus, 75
Carex, 151, 587; **agrostoides,** 607; australis, 151; **austrina,** 151, 152; **brevisquama,** 152; filiformis, 587; foetida, 606; Gayana, 603–605; Gayana hyalina, 603; glaucescens, 365; Goodenovii, 587; Hookeriana, 153–155; hyalina, 603; incurva, 606; **latebrosa,** 603, 604, 607; marcida, 603–605; Muhlenbergii, 152; Muhlenbergii australis, 151, 152; Muhlenbergii austrina, 151; **neomexicana,** 153, 154; occidentalis, 153, 154; **perglobosa,** 606; **simulata,** 603–605; spectabilis, 587; stenophylla, 605; **tumulicola,** 154, 155; vallicola, 153; vernacula, 606; verrucosa, 587; vulgaris, 587
Carex, Notes on, 151, 603
Carolinas and Virginia, A midsummer journey through the coastal plain of the, 351
Carpinites microphyllus, 193
Cassia, 230; tomentosa, 239
Castalia odorata, 183, 364
Castanea dentata, 306
Castilleja **ampliflora,** 39; **arcuata,** 35; Bradburyi, 39; cognata, 36; desertorum, 37, 41; fasciculata, 41; **gracillima,** 39; hispida, 39; **humilis,** 37; lanceolata, 37, 38; lauta, 37; **Leonardi,** 36; linariaefolia, 36; lutea, 40; lutescens, 37, 41; **magna,** 36; miniata, 38; occidentalis, 40; pallida, 40; **parvula,** 40; **Pecten,** 41; **pulchella,** 40, 41; **purpurascens,** 38; rhexifolia, 36, 37; rupicola, 39; Suksdorfii, 36; Tweedyi, 38; **variabilis,** 37; **viscida,** 38; **Vreelandii,** 38
Castasetum maculatum, 120; maculatum luteopurpuratum, 120
Caudalejeunea, 533, 553, 557; Cresentiae, 554, 556; Haenkeana, 554; harpaphylla, 554; **Lehmanniana,** 554, 557, 566, 567; Leiboldii, 554, 556, 557
Caulanthus, 428, 436; crassicaulis, 436; hastatus, 436
Celastrophyllum crenatum, 197, 206; Newberryanum, 197; undulatum, 197
Celtis occidentalis, 416
Centaurea, 405
Ceratolejeunea, 8, 10, 15, 17, 22; Breutelii, 13; brevinerivis, 7, 15; **portoricensis,** 15, 17, 33; valida, 15
Ceratophyllum submersum, 183
Cercis canadensis, 364
Ceriops Candolleana, 272
Ceuthospora, 312; **lunata,** 312
Chaetochloa italica, 396; macrosperma, 586; verticillata, 397; viridis, 396
Chamaebatia **australis,** 263; foliosa australis, 263
Chamaecyparis thyoides, 377
Cheiranthus, 68; asper, 66; pacificum, 66
Chenopodium, 82; rubrum, 82
Chloris elegans, 207, 397
Chlorocrambe, 435; **hastata,** 436
Chloroplegma, 508
Chondrophora nudata, 365
Chondrophyllum Nordenskioeldii, 198, 206; orbiculatum, 198
Cinnamomum, 200; Heerii, 188, 200, 206; intermedium, 200
Cissites acuminatus, 200; affinis, 200; Brownii, 200; harkerianus, 200; Heerii, 200
Citrullus vulgaris, 414
Cladium mariscoides, 581
Cladocephalus, 513; scoparius, 513
Cladosporium **Oxycocci,** 306
Cladothrix lanuginosa, 79
Clavaria amethystina, 102; **amethystinoides,** 102; Schaefferi, 102
Claytonia linearis, 317
Clethra alnifolia, 367
Cliftonia, 352, 358; monophylla, 361
Clinterium, 309, 310; obturatum, 309
Clitocybe **nobilis,** 97

Closterium Brebissonii, 599; granulatum, 600; laeve, 600
Coastal plain of the Carolinas and Virginia, A midsummer journey through the, 351
Cobaea, 226, 230; scandens, 239
Coccinia indica, 414
Code of Botanical Nomenclature, American, 167
Collybia dryophila, 345; **hirticeps,** 98; **subsulphurea,** 345; zonata, 98
Colocasia antiquorum, 390
Cololejeunea, 3, 5, 8; floccosa, 12
Colorado, Description of new fossil moss from, 139
Coltricia cinnamomea, 465; perennis, 466
Colura, 533
Comptonia vindobonensis, 192
Concerning some West American fungi, 207
Coniosporium, 84
Coniothyrium, 593
Contributions to the Mesozoic flora of the Atlantic coastal plain, 185
Convolvulaceae, Studies in the North American, 143
Convolvulus aquaticus, 149; erianthus, 149; humistratus, 147, 148; incanus, 70; Pickeringii, 149; sepium, 71; Sherardi, 148; tenellus, 148; trichosanthes, 148
Cook, M. T. The embryology of Rhizophora Mangle, 271; The embryology of Rhytidophyllum, 179
Corallina discoidea, 495; incrassata, 501; Flabellum, 495; tridens, 501
Coriolopsis **aneba,** 466; **badia,** 466; **dermatodes,** 466; occidentalis, 466; **phocina,** 466; **semilaccata,** 466
Coriolus, 468; abietinus, 467; atypus, 467; **cuneatiformis,** 467; elongatus, 467; **maximus,** 467; **murinus,** 467; **vernicipes,** 468
Cornus florida, 366
Coronopus, 68
Cortinarius, 86
Corydalis Brandegei, 426
Corymbis flava, 114
Costa Rican orchids, 113
Crataegus Cerronis, 462; punctata, 311; saligna, 462
Credneria, 199, 200; denticulata, 199; Zenkeri, 199
Cressa, 143
Crucianella, 228
Cucumis angulatus, 414; Anguria, 415; Melo, 414; odoratissimus, 414; sativus, 333, 415
Cucurbita, 222, 342; maxima, 415; moschata, 225; ovifera, 415; Pepo, 222, 225, 415
Cucurbitaceae, Some features of pollen-formation in the, 221
Cucurbitaria, 349; **erratica,** 349; Sorbi, 349
Cushman, J. A. A synopsis of the New England species of Tetmemorus, 599
Cyathula lappulacea, 79
Cyclanthera, 222, 224, 225, 237, 241, 334; explodens, 223; Hystrix, 406; pedata, 222
Cyclodictyon **cubense,** 574; limbatum, 574; Regnellii, 574
Cyclomycetella, 468
Cycloporellus, 468; **cichoriaceus,** 468; **microcyclus,** 468
Cycnoches Rossianum, 121; Warscewiczii, 121
Cynodon dactylon, 208
Cyperus Iria, 364; rotundus, 364
Cyrilla, 352; racemiflora, 365, 370
Cystispora endophylla, 312
Cystopus, 63; Amaranthacearum, 77; Amaranthi, 77; argentinus, 71, 72; Bliti, 77; brasiliensis, 73; candidus, 64, 69; Convolvulacearum, 69; Convolvulacearum Swertiae, 72; cubicus, 73; Cyathulae, 77; Euphorbiae, 84; Ipomoeae-panduranae, 69; Lepigoni. 71; Mikaniae, 73, 74; platensis, 79; Portulaceae, 83; pulverulentus, 73, 74; Salsolae, 84; Schlechteri, 84; sibiricus, 68; spinulosus, 73, 74; Tillaeae. 76; Tragopogonis, 69, 74; Tragopogonis spinulosus, 73; tropicus, 68

Daedalea amanitoides, 480; applanata, 480; Hobsoni, 480; Palisoti, 480; repanda, 480; sanguinea, 468
Dendroalsia abietina, 12
Dendrolejeunea, 552
Dentaria diphylla, 65, 66; laciniata, 66
Dewalquea groenlandica, 194
Dichromena latifolia, 365
Dicranella Perrottetii, 569
Dicranodontium **setosum,** 570
Dicranolejeunea, 4
Dicranopteris, 243–247, 250; **arachnoides,** 249; **Bancroftii,** 251-253, 258; **bicolor,** 251, 252; **Brunei,** 251, 253; **costaricensis,** 251, 253; **cubensis,** 251, 253; dichotoma, 257; **farinosa,** 252, 254; **flexuosa,** 251, 254; **fulva,** 252, 255, 256; **furcata,** 252, 257; **gigantea,** 249; **glabra,** 249; **glauca,** 249; **intermedia,** 252, 258; **jamaicensis,** 251, 258; **linearis,** 250, 255; **lon-**

gissima, 249; **mellifera,** 251, 259; **orthoclada,** 251, 259, 261; **palmata,** 252, 259; **pectinata,** 247, 250, 251, 260; **pteridella,** 251, 260; **retroflexa,** 251, 260; **strictissima,** 252, 261
Dicranostyles, 143; scandens, 143
Dicranum frigidum, 569; speciosum, 569
Dicrypta iridifolia, 121
Dictyogramme ensiformis, 449; lanceolata, 451
Diholcos Haydenianus, 420; **micranthus,** 420
Dinema paleaceum, 118
Diospyros primaeva, 188, 204
Diplasiolejeunea, 3
Diplopterygium, 249, 251
Dothiorella **aberrans,** 103; Asiminae, 103
Draba caroliniana, 68
Drepanolejeunea, 22
Dufourea, 144; glabra, 145; sericea, 144

Earliella **corrugata,** 468
Ecballium, 337
Eccilia Acus, 100; **cinericola,** 347; **Subacus,** 100; **unicolor,** 99
Echinochloa colona, 364
Echinospermum Lappula, 68
EDGERTON, C. W. Notes on a parasitic Gnomonia, 593
Eleocharis Robbinsii, 581
Elfvingia **Elmeri,** 476; tornata, 476
Elleanthus **caricoides,** 119, 124
Embryo of Zea Mays, An occurrence of glands in the, 125
Embryology of Rhizophora Mangle, The, 271
Embryology of Rhytidophyllum, The, 179
Entodon macropodus, 573
Entoloma **deminutivum,** 99; **modestum,** 347; **murinum,** 98; Peckianum, 99; sericeum, 99
Epidendrum auritum, 118; cochleatum, 117; lineare, 116; ochraceum, 118; paleaceum, 118; Stamfordianum, 118
Epilobium, 401
Equisetum, 226, 231, 240, 398
Erechtites hieracifolia, 404
Erigeron annuus, 404; philadelphicus, 394, 404; ramosus, 394
Eriocaulon decangulare, 365; septangulare, 581
Erysibe sphaerica Caryophyllacearum, 71
Eucalyptus, 202; attenuata, 203; Geinitzi, 201–203, 206; **linearifolia,** 203; nervosa, 203
Euosmolejeunea, 20; trifaria, 32
Eupatorium, 404; ageratoides, 404; compositifolium, 364; purpureum, 404, 406; rotundifolium, 366
Euphorbia eriogonoides, 364, 370; maculata, 364, 370
Eurotia lanata, 211
EVANS, A. W. Hepaticae of Puerto Rico, 1, 533
Evolvulus, 143, 149; ovalifolius, 146; prostratus, 149; rotundifolius, 149
Excretions by roots, The production of deleterious, 279

Fagus americana, 364, 366, 372
Favolus tenuis, 469; Wightii, 469
Features of pollen-formation in the Cucurbitaceae, 221
Ferns, Amercian, 243
Fevillea, 223–225, 237, 241; cordifolia, 223
Ficus, 199; daphnogenoides, 188, 194, 206; inaequalis, 194, 206; Woolsoni, 194, 206
Flabellaria luteofusca, 513
Flammula **betulina,** 100
Flora of Southern California, Studies on the, 263
Florissant, New species of fossil moss from, 139
Fomes Cornu-bovis, 478; **luzonensis,** 476; melanoporoides, 478; **philippinensis,** 477; pullus, 479; pyrrhocreas, 478; semilaccatus, 466; semitostus, 477
Fontinalis, 140; pristina, 139, 142
Fossil mosses, American, 139
Fouquieria splendens, 210
Fradelia, 508; fuliginosa, 508
Franseria tenuifolia, 75
Frasera, 72
Fraxinus caroliniana, 364
Frullania, 2, 4, 7, 539, 558, 566; dichotoma, 558; diffusa, 558; filicina, 558, 563; spathulistipa, 558
Fuckelia, 313
Funalia **philippinensis,** 469; **versatilis,** 469
Funaria calvescens, 570; hygrometrica, 570
Fungi, Concerning some West American, 207
Fungi, New species of, 97, 305, 345
Further notes on Halimeda and Avrainvillea, 491

Gaertneria acanthocarpa, 75; discolor, 76
GAGER, C. S. An occurrence of glands in the embryo of Zea Mays, 125
Ganoderma amboinense, 477; **subtor-**

natum, 477; **Williamsianum,** 478
Genus Albugo, The, 61
Genus Androsace in New Mexico, The, 517
Genus Antrophyum, The, 445
Geranium carolinianum, 400; dissectum, 400; maculatum, 400; pusillum, 400; Richardsonii, 400; Robertianum, 400
Gibbera, 310; obturata, 310
Glands in the embryo of Zea Mays, An occurrence of, 125
Gleditsia aquatica, 364
Gleichenia, 243, 244, 247, 260; axialis, 258; Bancroftii, 252, 258; bicolor, 252; bifida, 249; brevipubis, 262; Brunei, 253; dichotoma, 248, 255; farinosa, 254; flexuosa, 254; furcata, 257, 260; gigantea, 249; glaucina, 261; intermedia, 258, 261; intermedia dissitifolia, 259; intermedia flexuosa, 260; Liebmanni, 261; linearis, 250; longissima, 248, 249; Matthewsii, 259; mellifera, 259; orthoclada, 259; palmata, 259; pectinata, 248, 260; pteridella, 260; pubescens, 248–250, 253, 260; retroflexa, 260; revoluta, 253; rigida, 255; strictissma, 261; trachyrhizoma, 262
Gleicheniaceae, Preliminary review of the North American, 243
Gloephyllum **edule,** 480
Gloeosporium, 593; **minus,** 315
Glomerella, 315; rufomaculans **Vaccinii,** 314, 316
Glottidium vesicarium, 364
Glyphomitrium, 141; **Cockerelleae,** 140, 142
Glyptostrobus Ungeri, 140
Gnaphalium purpureum, 404; spathulatum, 404
Gnomonia, 593–597; Rubi, 594–597; tetraspora Rubi, 596; veneta, 595
Gnomonia, Notes on a parasitic, 593
Gomontia polyrhiza, 507
Gonolobus, 411, 412
Gordonia Lasianthus, 365, 369
Gossypium, 230
Grayia polygaloides, 591; spinosa, 591
Griffiths, D. Concerning some West American fungi, 207
Guignardia Bidwellii, 316; **Vaccinii,** 316
Guillenia, 429, 430; longirostris, 430; rostrata, 429
Gutierrhezia **bracteata,** 265; divergens, 265
Gymnosporangium, 459; **Betheli,** 459, 462; clavipes, 461; **durum,**460; globosum, 461; **inconspicuum,** 461, 463; Libocedri, 463
Gymnosporangium and Roestelia, New western species of, 459
Gyrotheca tinctoria, 365

Habenaria, 585; blepharigottis, 365, 581; cristata, 365; gracilis, 585
Halimeda, 491, 493, 495, 496, 499; cuneata, 494, 498, 499; discoidea, 495–500, 514, 515; favulosa, 501, 502, 504; gracilis, 491, 500; incrassata, 501, 503; incrassata ovata, 503; Lessonii, 497; macroloba, 491, 496; Monile, 499, 501, 502, 504; Opuntia, 494, 499; papyracea, 495–497; platydisca, 491, 497, 500; scabra, 492, 494, 500, 502, 504; **simulans,** 503, 504, 516; tridens, 493, 494, 501–504, 515; Tuna, 491–500, 502, 504, 514, 515; Tuna f. platydisca, 497
Halimeda and Avrainvillea, Further notes on, 491
Hamosa **atrata,** 48; **atratiformis,** 48
Hapalopilus gilvus, 470; licnoides, 470; **subrubidus,** 470
Haplocladium microphyllum, 574
Haplosporella, 103
Harper, R. M. A midsummer journey through the coastal plain of the Carolinas and Virginia, 351
Haynes, C. C. Two new species of Aytonia from Jamaica, 57
Hecisopteris **minima,** 457, 458; pumila, 458; Werckleana, 458
Hedera, 198, 201; primordialis, 201, 206
Hedysarum Mackenzii, 425; **utahense,** 424
Helianthus, 404–406; annuus, 404; divaricatus, 404; doronicoides, 405; grosseserratus, 405; hirsutus, 405; Maximiliani, 405; occidentalis, 405; scaberrimus, 405; strumosus, 405; trachaelifolius, 405; tuberosus, 127, 135, 136, 405
Helicosporium **Tiliae,** 103
Heliocarpus, 402; americanus, 402
Helminthosporium **inaequale,** 307
Helvella, 102
Hemionitis, 445; brasiliana, 452; cajanensis, 453; gigantea, 457; lanceolata, 451; lineata, 450
Hepaticae of Puerto Rico, 1, 533
Herberta, 561, 562
Herpestis verticillaris, 76
Hesperidanthus, 431, 434; **linearifolius,** 434
Hesperis, 430; matronalis, 66
Heteropterygium, 250, 251
Heterothrix, 432, 435; **longifolia,** 435; **micrantha,** 435
Hexagona cervinoplumbea, 469; ciliata,

470; cucullata, 470; orbiculata, 469; Thwaitesii, 469; **vibecinoides,** 470; Wightii, 469
Hibiscus, militaris, 364, 369, 373
Hibiscus militaris, 364, 369, 373
Hollick, A. & Britton, E. G. American fossil mosses, with description of a new species from Florissant, Colorado, 139
Holomitrium **Maxoni,** 572; proliferum, 573; Wrightii, 573
Holopterygium, 249–251
Homalobus brachycarpus, 49; campestris, 49, 50, 417; camporum, 50, 417, 418; decurrens, 50, 418; dispar, 419; **divergens,** 50, 417, 418; hylophilus, 50; **humilis,** 417; junceus, 49; junciformis, 49; **microcarpus,** 417; **oblongifolius,** 50, 418; **paucijugus,** 418; simplicifolius, 49; **stipitatus,** 419; **strigulosus,** 419, 420; tenellus, 418–420; tenuifolius, 50; **uniflorus,** 49
Homalolejeunea, 544
Hookeriopsis gracilis, 571
Hordeum vulgare, 135
House, H. D. Studies in the North American Convolvulaceae, 143
Howe, M. A. Phycological studies, 491
Hydnum **Kauffmani,** 348; **sulcatipes,** 101
Hymenophyllum, 12; ciliatum, 11
Hypericum fasciculatum, 364
Hypnum, 140; Brownii, 139, 142; columbianum, 140, 142; Haydenii, 139, 140, 142
Hypocopra, 314

Ibervillea, 329–335, 339, 340; Sonorae, 329, 330, 338, 340, 343
Ibervillea Sonorae, The stem of, 329
Ilex glabra, 367
Impatiens, 403; aurea, 403; biflora, 403; Nolitangere, 403
Index to American botanical literature (1901–1905), 51; (1906), 105, 267, 575; (1907), 161, 213, 319, 379, 439, 483, 521, 609
Inocybe **decipientoides,** 100; decipiens, 101
Inonotus **Elmerianus,** 471
Inula salicina, 74
Iodanthus linearifolius, 430, 434; pinnatifidus, 66
Ipomoea Batatas, 70; carolina, 70; incarnata, 71; Jalapa, 71; lacunosa, 70; leptophylla, 70; mexicana, 70; pandurana, 71; pandurata, 71, 81; Pes-caprae, 71; simulans, 71; triloba, 71
Irpex maximus, 467
Isochilus linearis, 116
Iva, 405; ambrosiaefolia, 75; ciliata, 75; xanthiifolia, 395, 405

Jamaica, Two new species of Aytonia from, 57
Journey through the coastal plain of the Carolinas and Virginia, A midsummer, 351
Jubula, 4, 566
Juglandites, 192
Juglans cinerea, 284
Juncus biflorus, 374; effusus, 364; scirpoides compositus, 374
Jungermannia, 557; birotunda, 18; brachiata, 546; cucullata, 12; dilatata, 558; filicina, 557, 558, 563–565; filiformis, 18; geminiflora, 18; granulata, 539; incrassata, 540; Mackaii, 543; spathulistipa, 558; squamata, 5; subfusca, 26; Tamarisci, 558; transversalis, 534, 537; trinitensis, 566
Juniperus californica utahensis, 460, 461; communis, 140; scopulorum, 460; virginiana, 364

Kalmia, 204; Brittoniana, 204
Kawakamia, 393
Kentrophyta **minima,** 420; **tegetaria,** 421; Wolfii, 421
Kern, F. D. New western species of Gymnosporangium and Roestelia, 459
Kirkwood, J. E. Some features of pollen-formation in the Cucurbitaceae, 221
Knox, A. A. The stem of Ibervillea Sonorae, 329
Koeleria, 209

Lactarii from Vermont, Some, 85
Lactarius, 85–95; affinis, 90, 94; alpinus, 90, 94; **aspideoides,** 87, 93; aspideus, 87; **Bensleyae,** 87, 95; camphoratus, 90, 95; Chelidonium, 93; cilicioides, 94; cinereus, 90, 94; circellatus, 86, 90, 94; deceptivus, 86, 90, 95; deliciosus, 90, 93; distans, 95; flexuosus, 90, 95; fuliginosus, 91, 93; Gerardii, 91, 95; griseus, 88, 91, 95; hysginus, 91, 94; Indigo, 93; insulsus, 94; **isabellinus,** 88, 93; ligniotus, 91, 93; **minusculus,** 88, 94; **nitidus,** 89, 95; **oculatus,** 86, 89, 95; pallidus, 95; paludinellus, 91, 95; pargamenus, 94; parvus, 91, 94; piperatus, 86, 94; pyrogalus, 91, 94; regalis, 86; resimus, 86, 91, 93; rimosellus, 86, 91, 95; **rufulus,** 346; rufus, 94, 346; subdulcis, 86, 89–91, 95; subdulcis oculatus, 89; subpurpureus, 91, 93;

theiogalus, 92, 93; torminosus, 92, 93; trivialis, 94; turpis, 92, 94; uvidus, 92, 93, 102; vellereus, 92, 95; volemus, 92, 95; **xanthogalactus,** 346; zonarius, 94
Laelia acuminata, 118; peduncularis, 118; rubescens, 118
Lagenaria vulgaris, 415
Larix, 226–228, 231, 237, 239
Lathyrus **brachycalyx,** 425; decapetalus, 425
Laurophyllum, 189
Laurus, 188, 189
Lavatera, 230, 239
Lejeunea, 4, 18, 533, 558; barbiflora, 540; Bongardiana, 546, 548, 549; brachiata, 546, 548; Breutelii, 7, 8, 13; catenulata, 7, 8, 10–12; complicata, 546; Crescentiae, 554, 556, 557; eulopha, 22; filicina, 563; filiformis, 18; floccosa, 12; geminiflora, 18; granulata, 539; Guilleminiana, 546, 548, 549; Haenkeana, 554; harpaphylla, 553, 554, 556, 557; incrassata, 540, 542; Lehmanniana, 553, 554, 556; Leiboldii, 554; Mariei, 30, 32; Montagnei, 20; Muelleriana, 27, 29, 30; portoricensis, 8, 10, 15; Sagraeana, 30; seminervis, 7, 8; squamata, 5; subfusca, 26, 27, 30; taeniopsis, 539, 540; tenuifolia, 18; transversalis, 537; transversalis Hookeriana, 540; vincentina, 542
Lemna, 364; minor, 302
Lentinus, 104; ventricosus, 104
Lenzites applanata, 480; Palisoti, 480; pallida, 480; platypoda, 480; repanda, 480
Leotia chlorocephala, 103; **punctipes,** 102
Lepidium alyssoides, 427; Bourgeauanum, 428; **brachybotryum,** 427; campestre, 65, 68; **Crandallii,** 427; densiflorum, 66, 428; Eastwoodiae, 427; **Fletcheri,** 428; Jonesii, 427; montanum, 427; ruderale, 428; sativum, 68; scopulorum, 427; virginicum, 66
Lepigonum medium, 81
Lepiota cepaestipes, 97; **xylophila,** 97
Leptilon canadense, 394
Leptochloa mucronata, 208; viscida, 208
Leptodontium brevisetum, 572; **perannulatum,** 572
Leptothyrium **Oxycocci,** 311
Leskea microphylla, 574
Lilium, 228, 231; Catesbaei, 365
Limnobium Spongia, 364
Limnorchis stricta, 585
Limonium, 368; carolinianum, 368; Nashii, 368
Lindera, 189
Liquidambar styraciflua, 367
Liriodendron, 196, 197; **dubium,** 196, 206; Meekii, 196; oblongifolium, 196; primaevum, 196, 197; semialatum, 196; Tulipifera, 206, 367
Liriodendropsis, 196, 197; angustifolia, 196; simplex, 196
Literature, American botanical (1901–1905), 51; (1906), 105, 267, 575; (1907), 161, 213, 319, 379, 439, 483, 521, 609
Long Island, A new Utricularia from, 579
Lophodermium melaleucum, 309; Oxycocci, 309
Lopholejeunea, 1, 21–23, 29, 32, 535, 536, 550, 553, 557; Anderssonii, 21; dentistipula, 32; eulopha, 22, 32; fimbriata, 32; **Howei,** 30, 32, 34; Muelleriana, 24, 27, 29, 30, 32, 34; Sagraeana, 24, 26–30, 32, 33
Loxogramme, 445; lineata, 450
Ludwigia maritima, 368; suffruticosa, 364; virgata, 369
Lupinus amplus, 41, argenteus, 43, 45; argenteus argophyllus, 43; Bakeri, 44; brevicaulis, 45; canescens, 43; decumbens, 44; dichrous, 44; **flavicaulis,** 43; foliosus stenophyllus, 42; holosericeus, 43; laxiflorus, 41, 42; **laxispicatus,** 42; leucophyllus, 44; Lyallii, 45; **Macounii,** 42; **macrostachys,** 44; **marianus,** 41; oreophilus, 43; ornatus glabratus, 41; pseudoparviflorus, 41; pusillus, 45, 46; **roseolus,** 44; **rubens,** 45; **scaposus,** 45; Scheuberae, 42; **stenophyllus,** 42; **subulatus,** 43; tenellus, 42
Lycopersicon Lycopersicon, 392
Lycopodium, 139, 140; rupestre, 377
Lysiostyles, 143; scandens, 143

MACKENZIE, K. K. Notes on Carex, 151, 603
Macromitrium Husnoti, 573
Macrophoma **tiliacea,** 348
Macropodium, 429; laciniatum, 429; nivale, 429
Madia sativa, 405
Magnolia, 188; amplifolia, 195; Capellinii, 188, 195, 206; glauca, 367, 371; grandiflora, 364, 371; longifolia, 195; macrophylla, 196; **Newberryi,** 195, 206; obtusata, 195
Malapoenna, 189
Malvastrum densiflorum, 264, 265; **viscidum,** 264
Marasmius, 98
Marchantia polymorpha, 240

Marchesinia, 533, 543–546; Bongardiana, 546; brachiata, 546, 548, 549, 567; Guilleminiana, 546; Mackaii, 544, 545; robusta, 546
Marshallia graminifolia, 365
Mastigolejeunea, 533, 549, 550, 552, 557, 559; auriculata, 549, 551, 552; crispula, 550
Matricaria, 73; matricarioides, 75
Maxillaria iridifolia, 121; **Valenzuelana,** 121
Maximowiczia, 330, 334; Sonorae, 330
Melica imperfecta, 209
Melothria, 225, 237; scabra, 415
Mertensia, 243, 260; arachnoides, 249; farinosa, 254; flexuosa, 254; fulva, 255; furcata, 257; glabra, 249; glaucescens, 250; gleichenioides, 261; grandis, 257, 258; palmata, 259; pectinata, 260; pubescens, 255; rigida, 254; subtrisperma, 254
Mesosorus, 243
Mesozoic flora of the Atlantic coastal plain, 185
Micrampelis, 223, 225, 227–229, 231, 234, 236, 237, 241, 242; lobata, 223, 406, 415
Microchloa indica, 208
Microporellus **subdealbatus,** 471
Microporus Perula, 472
Microthamnium reptans, 571; thelistegum, 574; Tuerckheimii, 571
Midsummer journey through the coastal plain of the Carolinas and Virginia, 351
Momordica balsamina, 415; charantia, 415
Montia linearis, 317; minor, 317
Mosses, Fossil, 139
Mosses from tropical America, 569
Mukia scabrella, 415
MURRILL, W. A. Some Philippine Polyporaceae, 465
Myrica, 191, 192; acuta, 191; brookensis, 191; callicomaefolia, 192; carolinensis, 373, 374; cerifera, 367, 373, 374; **elegans,** 191, 206; Gale, 374; Newberryana, 191; praecox, 191; pumila, 365; raritensis, 191; rigida, 192
Myrtophyllum, 202; Geinitzi, 201, 202; Warderi, 202, 203
Myrtus caryophyllata, 1

NASH, G. V. Costa Rican orchids, 113
Naucoria semiorbicularis, 101; **sororia,** 101; tabacina **bicolor,** 347
Neckera abietina, 11, 12; undulata, 573
Neottia flava, 114
Neslia paniculata, 67
Neurolejeunea, 1, 3, 5, 7, 8, 10, 15, 17; **Breutelii,** 13, 15, 33; catenulata, 9, 10, 12, 13, 15, 17, 33; portoricensis, 15; seminervis, 15
New England species of Tetmemorus, 599
New Mexico, The genus Androsace in, 517
New species of Aytonia from Jamaica, 57
New species of fossil moss from Florissant, 139
New species of fungi, 97, 305, 345
New species of Uredineae, 583
New Utricularia from Long Island, 579
New western species of Gymnosporangium and Roestelia, 459
New willows from the Canadian Rocky Mountains, 157
Nigrofomes melanoporus, 478
Nigroporus **durus,** 471; vinosus, 471
Nitrophila, 210; occidentalis, 210
Nomenclature, American Code of Botanical, 167
North American Convolvulaceae, 143
North American Gleicheniaceae, 243
North American Peronosporales, 61, 387
North Carolina, Contributions to the Mesozoic flora of, 185
Notarisia, 141
Notes on Carex, 151, 603
Notes on Halimeda and Avrainvillea, 491
Notes on a parasitic Gnomonia, 593
Nymphaea advena, 183; fluviatilis, 371
Nyssa biflora, 352, 367; Ogeche, 358; sylvatica, 352; uniflora, 367

Occurrence of glands in the embryo of Zea Mays, 125
Octoblepharum albidum, 570
Odontolejeunea, 2, 5, 536, 537
Oidium, 408
Omphalanthus, 1, 17, 18, 20; filiformis, 18, 20, 21, 33; geminiflorus, 18, 21
Omphalia, 98; **curvipes,** 345; grisea, 98; integrella, 345; pusillissima, 345; **serotina,** 98; **vestita,** 345
Omphalolejeunea, 18
Onoclea polypodioides, 243
Orchids, Costa Rican, 113
Ornithocephalus, 123
Orobus dispar, 419
Oxalis Wrightii, 264
Oxydendrum arboreum, 366
Oxygraphis Cymbalaria, 584
Oxypolis filiformis, 364
Oxytropis argentata, 424; Besseyi, 422, 424; monticola, 424
Ozonium auricomum, 305; **omnivorum,** 305

Pachypodium, 429, 430; integrifolium, 429, 433; laciniatum, 429; linearifolium, 434; sagittatum, 429
Panicum caespitosum, 209; digitarioides, 364; hebotes, 586; molle, 586
Parasitic Gnomonia, Notes on a, 593
Parietaria pennsylvanica, 407
Parthenium, 73; integrifolium, 75; repens, 75
Parthenocissus quinquefolia, 408; tricuspidata, 409
Passiflora incarnata, 364
PECK, C. H. New species of fungi, 97, 345
Peckiella hymenii, 102; **hymenioides,** 102
Pedicularis bracteosa, 35; Canbyi, 35; **siifolia,** 35
Peltandra undulata, 239
Peltolejeunea, 20
Penicillus, 513
Peperomia pellucida, 68, 81
Peronoplasmopara, 412, 413; Celtidis, 415; cubensis, 413
Peronospora, 395, 398, 399, 412, 413; australis, 406; Cactorum, 390; Celtidis, 415; conglomerata, 400; Conii, 401; cubensis, 412, 413; entospora, 393; Epilobii, 401; Geranii, 399; Gonolobi, 411; graminicola, 396; graminicola Setariae-italicae, 396; Halstedii, 403; Halstedii Ambrosiae, 403; illinoensis, 407; Impatientis, 402, 403; infestans, 389, 391; Kellermanii, 394; macrocarpa, 401; macrospora, 401; nivea, 401; nivea Geranii, 399; obducens, 402; ribicola, 410; sicyicola, 406; simplex, 394; Umbelliferarum, 401; viticola, 398, 407
Peronosporales, North American, 61, 387
Petalostemon, 589; candidus, 589; multiflorus, 589; oligophyllus, 589; purpureus, 589; villosus, 589; violaceus, 589
Phacidium, 308
Phalaris arundinacea, 398
Pharbitis hederacea, 71; purpurea, 71
Phaseolus lunatus, 390
Philippine Polyporaceae, 465
Phoenix dactylifera, 136
Phoma cystisporea, 312; leptidea, 308
Phoradendron flavescens, 367
Phragmicoma, 543; Bongardiana, 546, 548, 567; catenulata, 10, 12; catenulata incompleta, 12; filiformis, 18; granulata, 539; Guilleminiana, 546, 548; Haenkeana, 554, 556, 557; ocellulata, 5: transversalis, 537
Phragmites, 190, 191, 206
Phragmolejeunea, 552
Phragmopyxis, 590
Phycological studies, 491
Phyllosticta **putrefaciens,** 307
Phytophthora, 388, 389, 391; Cactorum, 389, 390; Colocasiae, 389, 390; infestans, 389, 391, 392, 408; Nicotianae, 389, 390; omnivora, 391; Phaseoli, 389, 390; **Thalictri,** 389, 392
Pieris floribunda, 352; nitida, 352
Pinckneya, 358
Pinus, 284; echinata, 366, 367; Elliottii, 360–362, 364, 373, 375; insularis, 481; palustris, 351, 360, 361, 375, 376; rigida, 376; serotina, 351, 362, 367, 376; Taeda, 358, 360, 361, 367, 375, 376; virginiana, 313
Pisum, 231
Plagiochasma elongatum, 58; Wrightii, 59
Plagiorhabdus, 310; **Crataegi,** 310, 311; **Oxycocci,** 311
Planera, 193, 373; aquatica, 364, 372; betuloides, 193; **cretacea,** 193, 206; Ungeri, 193
Plasmopara, 394, 395, 398, 399, 412, 413; australis, 406; Celtidis, 415; cubensis, 413; entospora, 394, 395; Epilobii, 401; Geranii, 399; Gonolobi, 411; Halstedii, 403; Heliocarpi, 402; Impatientis, 402; Kellermanii, 394; nivea, 401; obducens, 402; pusilla, 400; ribicola, 410; Viburni, 410; Vincetoxici, 411, 412; viticola, 408
Platanus, 373; affinis, 200; Heerii, 200; occidentalis, 364, 370, 372
Platylejeunea, 534; granulata, 539; transversalis, 537; vincentina, 542
Platyzoma, 243, 244
Pleurophragma, 431, 433; **gracilipes,** 433; **integrifolium,** 433, 434; **platypodum,** 434
Pleurothallis minax, 115, 116; plumosa, 115, 116; polyliria, 115; Valenzuelana, 121, 122
Poa, 584; arida, 584; nevadensis, 584
Poacites, 190
Pogonatum tortile, 571
Pollen-formation in the Cucurbitaceae, 221
Polygala cymosa, 364; lutea, 365, 370; ramosa, 365
Polymnia Uvedalia, 364
Polypodium dichotomum, 243, 250, 255; furcatum, 257; glaucum, 249; lineare, 250, 255
Polyporaceae, Some Philippine, 465
Polyporus affinis, 472; albomarginatus, 478; anebus, 466; atypus, 467; aureomarginatus, 470; australis, 476;

bicolor, 466; bivalvis, 469; breviporus, 470; brunneolus, 467; caliginosus, 478; carneoniger, 472; celebicus, 471; cervino-gilvus, 466; cinereofuscus, 478; **coracinus,** 471; corrugatus, 468; dermatodes, 466; discipes, 470; durus, 471; elongatus, 467; endotheius, 478; fastuosus, 479; fuscobadius, 468; grammocephalus, 472; holosclerus, 470; incomptus, 472; intybaceus, 468; iodinus, 468; Kermes, 478; laeticolor, 478; lanatus, 466; mariannus, 468; Meyenii, 467; microcyclus, 468; murinus, 467; ostreaeformis, 474; **palensis,** 472; pectinatus, 479; Peradeniae, 466; Perula, 472; phocinus, 466; polychrous, 472; scabrosus, 468; scorteus, 466; setiporus, 468; spadiceus, 470; surinamensis, 473; tornatus, 476; vernicipes, 468; vibecinoides, 470; vibecinus, 473; Wightii, 469; xanthopus, 472; zonalis, 473; zonalis semilaccatus, 466
Polypremum procumbens, 364
Polystictus badius, 466; cichoriaceus, 468; cilicioides, 470; Didrichsenii, 467; Persoonii, 468
Polytaenium, 447–449; lanceolatum, 450; lineatum, 450
Polytrichadelphus ciliatus, 571
Ponera amethystina, 116
Pontederia cordata, 364
Populites affinis, 200
Populus 364, 373; deltoides, 373
Porana, 143; velutina, 144
Portulaca oleracea, 81, 83
Potentilla, 284; fruticosa, 284
Preliminary review of the North American Gleicheniaceae, 243
Prevostea, 143, 144; amazonica, 145; ferruginea, 145; glabra, 145; sericea, 144, 145; spectabilis, 146; umbellata, 146; velutina, 144
Production of deleterious excretions by roots, 279
Prospodium Amphilophii, 588; appendiculatum, 588; **bahamense,** 587
Protomyces graminicola, 396
Prunus, 284
Psathyrella **betulina,** 101; **caespitosa,** 348
Pseudoperonospora, 389, 399, 412, 413; **Celtidis,** 407, 413, 415; cubensis, 406, 413
Pseudoplasmopara, 413
Pseudotsuga Douglasii, 591; mucronata, 591
Pteridium aquilinum, 367
Pteris dichotoma, 243, 257
Pterocaulon undulatum, 365
Pterospermites, 188, 199; **carolinensis,** 198, 206; cordifolius, 199; **credneriifolius,** 199, 206; modestus, 199
Ptychanthus, 4, 566
Ptychocoleus, 543
Ptychomitrium, 141
Puccinia, 585, 587; Agrostidis, 584; Caricis, 591; **Chaetochloae,** 585; **cinerea,** 583; coronata, 585; epiphylla, 584; eslavensis, 586; **Eurotiae,** 210; Gonolobi, 411; minuta, 587; **minutissima,** 587; **panicicola,** 586; **perminuta,** 584; Petalostemonis, 589; Poarum, 584; **praegracilis,** 585; **quadriporula,** 586; Rhamni, 585
Puerto Rico, Hepaticae of, 1, 533
Pycnoporus sanguineus, 473
Pyropolyporus **albomarginatus,** 478; **caliginosus,** 478; **endotheius,** 478; **fastuosus,** 479; **lamaensis,** 479; **Merrillii,** 479; **pectinatus,** 479; **Williamsii,** 479

Quamoclit Quamoclit, 71
Quercophyllum, 193
Quercus, 188, 192, 193; alba, 366; Catesbaei, 351, 367, 372; cinerea, 365; coccinea, 306; digitata, 367; Ellsworthiana, 192; geminata, 361; Johnstrupi, 192, 193; Marioni, 192; marylandica, 367; minor, 366; Phellos, 366, 367; **Pratti,** 192, 206; sphenobasis, 192; virginiana, 364; Wardiana, 192

Raphanus sativus, 65, 67, 81
Ravenelia, 592
Reboulia hemisphaerica, 58
Reed, H. S. & Schreiner, O. The production of deleterious excretions by roots, 279
Reinwardtia, 144; glabra, 145; sericea, 145
Reseda, 65
Review of the North American Gleicheniaceae, 243
Rhabdospora **Oxycocci,** 311
Rhexia Alifanus, 365
Rhinotrichum **Sumstinei,** 103
Rhipilia longicaulis, 508–510; Rawsoni, 510; tomentosa, 512
Rhizogonium spiniforme, 571, 573
Rhizophora Mangle, 271; mucronata, 272
Rhizophora Mangle, The embryology of, 271
Rhizopus, 285

Rhus, 192; acuminata, 192; copallina, 366
Rhynchostegium conchophyllum, 571; Knowltoni, 140, 142
Rhysotheca, 389, 398, 399, 413; **australis,** 399, 406, 407; **Epilobii,** 399, 401; **Geranii,** 399; **Gonolobi,** 399, 411; **Halstedii,** 399, 403; **Heliocarpi,** 399, 402; **illinoensis,** 399, 407; **obducens,** 399, 402; **ribicola,** 399, 410; **Umbelliferarum,** 399, 401; **Viburni,** 399, 410; **viticola,** 399, 407
Rhytidophyllum, 179; crenulatum, 179, 183, 184; tomentosum, 179, 183, 184
Rhytidophyllum, The embryology of, 179
Ribes albinervum, 411; divaricatum, 411; hirtellum, 411; oxycanthoides, 411; prostratum, 411; rotundifolium, 411; rubrum, 411; rubrum subglandulosum, 411
Ricinus, 128; communis, 135
Rigidoporus **surinamensis,** 473
Rocky Mountain flora, Studies on the, 35, 417
Rocky Mountains, Two new willows from the Canadian, 157
Roestelia, 459; **Betheli,** 461; **Harknessiana,** 462, 463; **Harknessianoides,** 463
Roestelia, New western species of, 459
Roots, The production of deleterious excretions by, 279
Roripa Armoracia, 65, 67; hispida, 67; Nasturtium, 68; obtusa, 67; palustris, 67; sessiliflora, 67; Walteri, 67
Rowlee, W. W. Two new willows from the Canadian Rocky Mountains, 157
Rubus nigrobaccus, 593, 594
Rudbeckia fulgida, 394; laciniata, 405; triloba, 405
Russula, 86
Rydberg, P. A. Studies on the Rocky Mountain flora, 35, 417
Rynchospora corniculata, 364; fusca, 581; semiplumosa, 375

Sabal glabra, 364
Sabbatia campanulata, 364; lanceolata, 365
Sabina scopulorum, 460; utahensis, 460, 461
Sagittaria variabilis, 183
Salix, 157, 189; **albertana,** 157, 158; Barrattiana, 158; Barrattiana angustifolia, 158; glaucops, 159; lucida, 159; **Maccalliana,** 158; nigra, 367; Tweedyi, 158; wyomingensis, 157
Sambucus canadensis, 364
Sarcobatus, 210
Sarracenia flava, 365, 371; minor, 365
Sassafras, 201; acutilobum, 201; harkerianum, 200; hastatum, 201; variifolium, 369
Saururus cernuus, 183
Schoenocrambe linifolium, 67
Schomburgkia, 118
Schreiner, O. & Reed, H. S. The production of deleterious excretions by roots, 279
Sclerospora, 388, 395; **Farlowii,** 207, 396, 397; graminicola, 396, 397; Kriegeriana, 398; macrospora, 398; Magnusiana, 398
Scoliosorus, 447–450; ensiformis, 449
Secale, 221
Selaginella acanthonota, 377
Sematophyllum admistum, 574; subsimplex, 571
Senecio, 73; aureus, 76; cymbalaroides, 75; Hartianus, 75; lugens, 76; oblanceolatus, 76; peninsularis, 76; serra, 76; serra integriusculus, 76; tomentosus, 367
Septoria **longispora,** 308
Sequoia heterophylla, 188, 189
Serapias flava, 114
Serenoa, 358; serrulata, 361
Shear, C. L. New species of fungi, 305
Sicydium, 330; Lindheimeri, 330
Sicyos angulatus, 406, 415
Sida rhombifolia, 364
Silphium, 221, 239; integrifolium, 405; laciniatum, 405; perfoliatum, 405; terebinthinaceum, 405; trifoliatum, 405
Sinapis alba, 68
Sisymbrium officinale, 67
Smilax lanceolata, 364; laurifolia, 365
Solanum tuberosum, 392
Solidago canadensis, 405; Riddellii, 405; rigida, 394
Some features of pollen-formation in the Cucurbitaceae, 221
Some Lactarii from Vermont, 85
Some Philippine Polyporaceae, 465
Some West American fungi, 207
Sophia andrenarum, 209; brachycarpa, 436; filipes, 436; Hartwegiana, 67; incisa, 67, 436; intermedia, 436; **magna,** 436; millefolia, 67; **Nelsonii,** 436; pinnata, 65, 67, 437
Sordaria, 314
Sorosporium Eriochloae, 209; Montiae, 317; **ovarium,** 209
Southern California, Studies on the flora of, 263

Spartina glabra, 367
Species of Antrophyum, American, 445
Species of Aytonia from Jamaica, 57
Species of fungi, New, 97, 305, 345
Species of Gymnosporangium and Roestelia, New western, 459
Species of Uredineae, New, 583
Sphaeria obturata, 309, 310; obturata epiphylla, 309; obtusata, 309; Sclerotium, 310
Sphaeronema **pomorum,** 307
Sphaeropsis, 309; Asiminae, 103
Sphagnum, 92; medium purpurascens, 569
Spongipellis **luzonensis,** 473
Sporonema, 310; **epiphyllum,** 309; obturatum epiphyllum, 309; **Oxycocci,** 308; phacidioides, 310; **pulvinatum,** 308
Sporotrichum **Quercuum,** 306; sulphureum, 306; sulphureum f. Quercuum, 306
Standley, P. & Wooton, E. O. The genus Androsace in New Mexico, 517
Stanfordia, 428
Stanleya, 431
Stanleyella, 431, 435; **Wrightii,** 435
Statice caroliniana, 368; Limonium, 368
Stem of Ibervillea Sonorae, 329
Sterculia Saportanea, 200
Stictolejeunea, 1–5, 9, 10, 536, 545, 560; Kunzeana, 1, 3, 7; squamata, 1–3, 5, 7, 13, 33
Streptanthus, 428, 430, 431; linearifolius, 430, 434; longifolius, 435; longirostris, 430; maculatus, 431; micranthus, 435; wyomingensis, 431, 432
Streptochaeta, 126
Stromatopteris, 243, 244
Studies, Phycological, 491
Studies in North American Convolvulaceae, 143
Studies in North American Peronosporales, 61, 387
Studies on the flora of Southern California, 263
Studies on the Rocky Mountain flora, 35, 417
Stylisma, 143, 147, 149; **angustifolia,** 147, 148; aquatica, 148, 149; elliptica, 149; evolvuloides, 148; evolvuloides angustifolia, 149; humistrata, 147, 148; peduncularis, 147, 148; Pickeringii, 148, 149; **rotundifolia,** 148, 149; **trichosanthes,** 147, 148; **villosa,** 148, 149
Swertia, 72; connata, 72, 81
Symbiezidium, 533–535, 537, 544, 545, 550; **barbiflorum,** 540, 542, 567; Breutelii, 13; catenulatum, 10; granulatum, 539, 540; squamatum, 5; transversale, 534, 537, 538, 540–543, 567; vincentinum, 542, 543
Symplocarpus, 228; foetidus, 239
Synopsis of the New England species of Tetmemorus, 599
Syrrhopodon Hobsoni, 573

Taxilejeunea, 18
Taxodium distichum, 352, 367; imbricarium, 364, 376
Tecoma bahamensis, 588; radicans, 364
Tetmemorus, 599–601; Brebissonii, 599, 601; granulatus, 600, 601; laevis, 600, 601; minutus, 600, 601
Tetmemorus, A synopsis of the New England species of, 599
Thalictrum, 234; purpurascens, 234, 392
Thelypodiopsis, 431, 432; **aurea,** 432; **Bakeri,** 432; **elegans,** 432, **wyomingensis,** 432
Thelypodium, 428–431; aureum, 430, 432; Bakeri, 430, 432; elegans, 430, 432; integrifolium, 430, 433, 434; integrifolium gracilipes, 433; laciniatum, 429, 433; lasiophyllum, 429, 430; **leptosepalum,** 432; lilacinum, 434; linearifolium, 430, 434; longifolium, 431, 435; micranthum, 431, 435; **Palmeri,** 432; sagittatum, 433; utahense, 429; Wrightii, 431, 434, 435
Thladiantha, 330, 337, 339, 340
Thlaspi Bursa-pastoris, 68; glaucum, 67; Nuttallii, 67
Thyella tamnifolia, 71
Thysananthus, 552, 557, 558, 560, 566
Tilia americana, 103, 348
Tillaea rubescens, 76
Tillandsia usneoides, 364, 375
Timmiella subanomala, 571
Tissa leucantha, 72; marina, 72
Tium sparsiflorum, 48; **variegatum,** 47
Tofieldia racemosa, 365
Torula, 84
Tragopogon porrifolius, 76, 81
Trametes Aurora, 467; badia, 466; bicolor, 468; **caespitosa,** 473; colliculosa, 480; dermatodes, 466; **lamaensis,** 474; **luzonensis,** 474; Muelleri, 474; nitida, 468; **ostreaeformis,** 474; **subacuta,** 474; versatilis, 469; **Williamsii,** 475
Trichantha, 146
Trichodium, 584
Trichosanthes colubrina, 415

Trichostema lanatum, 265; lanatum denudatum, 265; Parishii, 265
Trifolium **Aitonii,** 46; **confusum,** 46; dasyphyllum, 47; **inaequale,** 47; latifolium, 46; longipes, 46; montanense, 47; Parryi, 47; pedunculatum, 46; Rusbyi, 46; Rydbergii, 46; **uintense,** 47
Triticum, 221; vulgare, 286, 398
Tropical America, Mosses from, 569
Tsuga canadensis, 583, 591; heterophylla, 583; Mertensiana, 583
Two new species of Aytonia from Jamaica, 57
Two new willows from the Canadian Rocky Mountains, 157
Typha latifolia, 364
Tyromyces **Elmeri,** 475

Udotea, 495, 512, 513; conglutinata, 513; Desfontainei, 513; flabellata, 495; Flabellum, 495; luteofusca, 513; tomentosa, 512
Underwood, L. M. American ferns, 243
Uredineae, New species of, 583
Uredo, 63; Bliti, 77; candida, 63, 64; candida Tragopogi, 72; Chaetochloae, 585; Cheiranthi, 64; Convolvulae, 68; **Grayiae,** 591; Holwayi, 583; **inquirenda,** 592; Portulacae, 83; Tragopogi, 72
Urocystis Agropyri, 209; **Sophiae,** 209
Uropyxis, 589, 590; Amorphae, 590; Petalostemonis, 589, 590
Ustilago Boutelouae, 208; Calandriniae, 317; **Claytoniae,** 317; Cynodontis, 208; heterogena, 208; hypodites, 208; **Microchloae,** 207
Utricularia, 579, 580; cornuta, 579; juncea, 579–581; resupinata, 581; simplex, 581; **virgatula,** 580–582
Utricularia from Long Island, A new, 579

Vaccinium crassifolium, 365; macrocarpum, 307–309, 311, 312, 314–316
Vaucheria geminata, 507; hamata, 507
Verbesina encelioides, 405
Vermont, Some Lactarii from, 85
Vernonia angustifolia, 365; Baldwinii, 405; noveboracensis, 405, 406
Viburnum acerifolium, 410; dentatum, 410; nudum, 410; Opulus, 410; pubescens, 410
Vigna Catjang, 287
Vignea, 131
Vincetoxicum gonocarpos, 412; **hirsutum,** 411, 412; suberosum, 412
Virginia, A midsummer journey through the coastal plain of, 351
Vitis, 404, 409; aestivalis, 408, 409; bicolor, 409; californica, 409; cinerea, 409; cordifolia, 409; labrusca, 409; riparia, 409; rotundifolia, 364, 409; vinifera, 409; vulpina, 409
Vittaria, 451, 457, 458; costata, 457; lanceolata, 450; lineata, 458

Warscewiczella Wendlandii discolor, 123
Wedelia incarnata, 80
West American fungi, 207
Western species of Gymnosporangium and Roestelia, New, 459
Widdringtonia helvetica, 140
Wilberforcia, 144
Williams, R. S. Mosses from tropical America, 569
Willows from the Canadian Rocky Mountains, New, 157
Wilson, G. W. Studies in North American Peronosporales, 61, 387
Windham County, Some Lactarii from, 85
Wooton, E. O. & Standley, P. The genus Androsace in New Mexico, 517

Xanthium canadense, 405
Xanthoxalis **californica,** 264; **Wrightii,** 264
Xylobium foveatum, 121
Xylophacos amphioxys, 49; **aragalloides,** 48
Xylothermia montana **tomentosa,** 263

Zea Mays, 125, 132, 136, 287
Zea Mays, An occurrence of glands in the embryo of, 125
Zostera, 221; marina, 240
Zygadenus glaberrimus, 365
Zygostates, 123; **costaricensis,** 122, 124